

REPORT DOCUMENTATION PAGE

AFRL-SR-BL-TR-02-

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be penalized for providing false or misleading information. PLEASE DO NOT RETURN YOUR FORM TO 1.

1. REPORT DATE (DD-MM-YYYY) 19 February 2002		2. REPORT TYPE Final Technical		3. DATES COVERED (From - To) 01/01/97 - 12/31/00	
4. TITLE AND SUBTITLE Long-Lived, Energetic States of Small Molecules: Spectroscopy, Pattern Recognition, and Formation/Destruction Mechanisms				5a. CONTRACT NUMBER 00-142-5594	
				5b. GRANT NUMBER F49620-97-1-0040	
				5c. PROGRAM ELEMENT NUMBER 611022F	
6. AUTHOR(S) Robert W. Field Robert J. Silbey				5d. PROJECT NUMBER	
				5e. TASK NUMBER 2303/ES	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Massachusetts Institute of Technology 77 Massachusetts Avenue Cambridge, MA 02139-4307				8. PERFORMING ORGANIZATION REPORT 20020315 073	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) AFOSR/NL 801 N RANDOLPH ST RM 732 ARLINGTON VA 22203-1977				10. SPONSOR/MONITOR'S ACRONYM(S) AFOSR/NL	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT DISTRIBUTION STATEMENT A Approved for Public Release Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Metastable, electronically excited states of atoms and small molecules have chemical and photophysical properties that are relevant to Air Force missions in communication, upper atmospheric modeling, and high speed vehicle detection, tracking, and identification. A second-generation supersonic molecular beam apparatus with optimized multispectral capabilities has been designed, constructed, and tested. A series of experiments on acetylene has defined the fundamental mechanisms of three complementary signal channels: UV-LIF (Ultraviolet Laser Induced Fluorescence), IR-LIF (Laser Induced Infrared Fluorescence from photofragments), and SEELEM (Surface Electron Ejection from Laser Excited Metastable molecules). Simultaneous measurements of S_1 , T_1 , and T_3 characters of each eigenstate permit the causal mechanisms of photoexcitation, intersystem crossing, and collisional deexcitation to be identified. Powerful statistical pattern-recognition techniques have been developed which permit the disentanglement of key structural and dynamical features in the spectra and the ability to distinguish between <i>direct</i> (statistical) and <i>doorway-mediated</i> (causal) limiting mechanisms for intersystem crossing.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF: Unclassified			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON Robert W. Field
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (include area code) 617-253-1489

FINAL TECHNICAL REPORT

Long-Lived, Energetic States of Small Molecules: Spectroscopy, Pattern Recognition, and Formation/Destruction Mechanisms

Air Force Office of Scientific Research
Grant Number F49620-97-1-0040

Co-Principal Investigators: Robert W. Field
Robert J. Silbey

Massachusetts Institute of Technology
Department of Chemistry
77 Massachusetts Avenue
Cambridge, Massachusetts 02139-4307

Period Covered: 1 January 97 - 31 December 2000

I. Staff Supported by AFOSR Grant

A. Postdocs

1. C. Gabe Morgan, currently postdoc, CalTech
2. Steve Drucker (1/97 - 7/98), currently Assistant Professor, Department of Chemistry, University of Wisconsin, Eau Claire
3. Manjula Canagaratna (9/99 - 6/00), currently staff scientist, Aerodyne Corporation

B. Graduate Students

1. Kevin Cunningham (1/97 - 6/00), currently staff scientist, Applied Materials
2. Selen Altunata (9/99 - 12/00), currently staff scientist, IGEN Corporation
3. Ryan Thom (6/00-12/00), MIT
4. Xing Jiang (6/99), MIT.

C. Undergraduate Students

1. Heather Drake (1/97)
2. Simon Tisminezky (9/97 - 8/97)
3. Matt Bonnet (7/97 - 8/97)
4. Seth Karon (1/99)
5. Kuo-Chang Lian
6. Susan Lang

II. Research Accomplishments (1 January 1997 – 31 December 2000)

The following is a slightly edited version of the "Accomplishments" section (and cited References) of the 3-year Renewal Proposal, submitted in February 2000, and currently supported under AFOSR F49620-01-1-0078.

The primary experimental apparatus constructed for this project is a two-chamber vacuum system which houses a pulsed, skimmed, supersonic molecular beam, equipped with three complementary detection schemes: a Surface Electron Ejection by Laser Excited Metastables (SEELEM) [SNE86ab, 88, 91] detector (with Time-of-Flight and interchangeable Au and Cs detector surface capabilities), a Germanium near-infrared (800-1700 nm) detector, and a photomultiplier (200-800 nm). This is a second-generation apparatus that was designed based on our experience with a much simpler single-chamber apparatus [HUM97]. The key feature of this apparatus is that four signal channels may be simultaneously recorded: parent molecule SEELEM (late TOF gate), metastable photofragment SEELEM (early TOF gate), infrared fluorescence excited by an ultraviolet laser (IR-LIF), and UV-LIF. This *multi-spectral* capability provides unique insights into the electronic character of the metastable species. These insights are based on a detailed and quantitative understanding of the specific characteristics of molecular eigenstates to which each detection channel is most sensitive, in other words, the "detection mechanism." In particular, each molecular eigenstate can be a mixture of several electronic states (excited singlet state, S_1 , lowest triplet state, T_1 , excited triplet state, T_i $i > 1$, and electronic ground state, S_0) and each signal channel has different detectivities for the fractional characters of each type of electronic state. Far from being spectroscopic arcanae, these fractional characters encode the chemical and photophysical properties from which fundamental excitation and decay mechanisms and practical quantitative measurements may be derived.

Any multispectral experiment will rapidly generate an enormous amount of spectral information. Advanced pattern recognition methods [COY95, 97, JAC97, 98] are essential for processing this deluge of

information. A suite of powerful pattern recognition schemes has been developed [ALT99, 00ab, CUN00b]. A key feature of these computer-automated, statistical schemes is that neither the nature nor the number of patterns being recovered from the raw multispectral data need be known in advance [JAC97, COY97]. One scheme, eXtended spectral AutoCorrelation (XAC), is designed to disentangle overlapping patterns associated with different values of the rigorously conserved upper state rotational quantum number, J' [CUN00b]. Each J' might fractionate into tens of resolved features and might overlap with the fractionated patterns belonging to several J' -values. Since each J' pattern is exactly replicated three times in a single vibronic band, via $R(J-1)$, $Q(J)$, and $P(J+1)$ rotational transitions, XAC is able to recover each J' pattern in a statistically rigorous manner. Once the J' -patterns are disentangled, another class of pattern recognition scheme is able to make a qualitative distinction between two extreme mechanisms of coupling between the well characterized *bright* state and the unknown manifold of *dark* states (*direct* and *doorway mediated* coupling) and to make quantitative determinations of the *dark* state molecular constants and the *bright* state-*dark* state coupling matrix elements [ALT00ab]. These computer-automated pattern recognition schemes are at least as important as the experimental apparatus in which the spectra are generated, because none of the spectra remotely resemble textbook spectra. The combination of a uniquely designed *multispectral* data set and a *statistical pattern recognition* approach provides the key to extracting information about a spectroscopic and dynamical problem of unprecedented complexity.

Although the experimental apparatus, experimental techniques, and statistical pattern-recognition methods described here are applicable to a wide range of small molecules, the following discussion applies specifically to the acetylene molecule, HCCH.


The discussion of our accomplishments during this project is organized as follows:

1. The Molecular Beam Apparatus.
2. The Mechanisms of the SEELEM and UV-LIF Detection Processes.
3. The Unexpectedly Sensitive Yet General Near Infrared Fluorescence Detection Channel.
4. Statistical Measures.
5. Dispersed Fluorescence Studies of Spectroscopic Perturbations in HCCH S_1 $3v_3$.
6. UV Cavity Ringdown.
7. Hg* Photosensitization Experiments.
8. Collaborations at AFRL, Hanscom Research Site: Pattern Recognition and Baseline Stripping.
9. The Spectroscopic Signature of Bond-Breaking Isomerization in HCP.

1. The Molecular Beam Apparatus.

The molecular beam portion of this project began with a collaborative experiment in the laboratory of Professor Alec Wodtke at the University of California, Santa Barbara [HUM97]. The UCSB apparatus was a sophisticated and complex apparatus designed for photofragment TOF spectroscopy and the scattering of highly excited species off surfaces [MOR96]. The features of the UCSB apparatus deemed essential for the initial concept-evaluation experiments planned at MIT were incorporated into the design of a first-generation single-chamber apparatus. All of the results presented in Sections 1-5 below were obtained using the single chamber apparatus. Based on those results, a second-generation two-chamber apparatus has been designed, assembled, and is currently being tested.

Figure 1. Schematic of two-chamber pulsed supersonic molecular beam apparatus. The excitation region, where the UV laser crosses the molecular beam, is viewed by a PMT (UV-LIF) and a Ge IR detector (IR-LIF). Laser excited metastable molecules travel at least 3 cm before impacting the Cs or Au metal surface of the SEELEM detector. The SEELEM signal channel has Time-Of-Flight capability, which can distinguish signals due to photons, parent metastables, and metastable photofragments.



The most important feature of the metastable molecular beam apparatus is the capability of equipping the SEELEM detector with clean, heated metal surfaces of low (Cesium, 2 eV) and high (Gold, 5 eV) work function and being able to switch from one detector surface to another without altering the detector location or active area. Owing to the extreme reactivity of Cs (and the 1×10^{-4} Torr operating pressure of our single-chamber apparatus), the Cs surface must be continuously vacuum deposited onto a rotating disc designed so that the deposition and detection functionalities are spatially separated. The SEELEM detector (a metal plate, electron optics, and electron multiplier) is sensitive exclusively to electronically excited molecules with vertical electronic excitation energy greater than the work function of the metal surface [HAG54]. It is well established that translational and vibrational energy does not contribute to the SEELEM detectivity [LIS80]. (Vibrational excitation *can* turn on SEELEM detectivity by giving Franck-Condon access to a nuclear geometry for which the vertical electronic excitation energy is larger than the work function of the metal.) When an electronically excited molecule impacts the metal surface, an electron is ejected, collected, multiplied, and detected.

The ability to switch between two metal surfaces, with high and low work functions, enables experimental determination of the fractional composition of each SEELEM-detectable eigenstate in terms of the energetically accessible electronic states: e.g. for acetylene, S_1 , T_3 , (T_1 , T_2) (abbreviated hereafter as T_1), and S_0 . Measurement of these electronic mixing fractions is equivalent to determining the excitation, unimolecular decay, and detection mechanisms. It was our expectation that S_1 and T_3 *but not* T_1 or S_0 fractional characters would be detectable on Au, and that S_1 , T_3 , and T_1 characters would be detectable on Cs. In this way we expected to be able to show that the “*bright*” S_1 state couples selectively to one special “*doorway*” triplet state, T_3 , and that this doorway state couples to a dense manifold of background “*dark*” triplet states, T_1 .




Figure 2. Level diagram for *Doorway Mediated* Intersystem Crossing. The T_3 doorway state promotes ISC by providing an indirect coupling pathway between the S_1 bright and the T_1 and S_0 dark states.

Furthermore, we expected to characterize this *doorway* state, thereby providing a causal (as opposed to phenomenological or statistical) mechanism for the InterSystem Crossing (ISC) [BIX67, AVO77] process in acetylene.

We discovered that, for the ~ 100 μ s transit times from the laser excitation region to the SEELEM detector surface, the Au and Cs surfaces are both primarily sensitive to S_1 and not T_3 character. This is a surprising result, with fundamental implications about the SEELEM detection process (both mechanism and quantum efficiency) in molecules (see Section 2, below) [SNE88]. Despite the unexpectedly low sensitivity of the SEELEM detector to triplet vs. excited singlet character, we were able to infer the fractional S_1 and T_3 characters in SEELEM-detectable eigenstates by exploiting the quite different dependence on S_1 character of the SEELEM vs. UV-LIF detection channels. This enabled us to demonstrate that ISC in HCCH S_1 is **Doorway Mediated** and to determine the properties of the *doorway* state.

We also know, by significantly increasing the transit time from excitation region to SEELEM detector surface, that it will be possible to determine the S_1 vs. T_3 relative sensitivities on Au and Cs surfaces and, by comparing SEELEM spectra recorded on Au and Cs, to measure directly the fractional T_3 and T_1 characters in the detected eigenstates. In order to compensate for the much higher sensitivity to S_1 than triplet character, it is necessary to ensure, by a long flight time from excitation to detector, that the only metastable molecules reaching the detector are those with fractional S_1 character a factor of at least 10 smaller than triplet character. Our second-generation apparatus has been designed with the possibility of varying the excitation to detector transit time from 20 μ s (3 cm in a He beam) to 1400 μ s (81 cm in an Ar beam), which is to be contrasted with minimum and maximum transit times of 80 and 200 μ s in our first-generation apparatus.

The first-generation apparatus is an unskimmed pulsed jet. The jet is crossed by the excitation laser 1 cm downstream from the nozzle. The excitation region is imaged ($f/2$ optics) onto a photomultiplier for detection of UV-LIF. Since near-IR-LIF turned out to be unexpectedly intense, we improvised an optical system ($\sim f/10$ optics) for imaging IR-LIF from the excitation region onto a liquid N_2 cooled Ge detector. The SEELEM detector is located ~ 12 cm downstream from the excitation region. The signal from the SEELEM detector consists of as many as three temporal features: UV fluorescence (and laser scatter) at ~ 0 delay, *metastable photofragments* at early time, and metastable parent molecules at later time [HUM97]. The carrier of the metastable photofragment signal is unknown (but most likely a >5 eV excited metastable state of CCH), but its formation from unexcited acetylene requires absorption of at least two 5.5 eV UV photons!

The SEELEM-TOF spectrum, in principle, contains information about the distribution of radiative lifetimes of the SEELEM-detectable eigenstates. However, this information is corrupted by two instrumental artifacts that cause a rapid decay, during each molecular beam pulse, in the intensity of the parent metastable feature of the TOF-SEELEM signal. These artifacts, which provide the primary motivation for redesign of the apparatus, are due to (i) attenuation and heating of the unskimmed molecular jet by the buildup of background gas scattered off the SEELEM detector surface, and (ii) reduced efficiency of the SEELEM surface during each gas pulse owing to inter-pulse reversible contamination by the enormous flux of molecules from the jet. There is, however, also the unevaluated effect on the SEELEM detector quantum efficiency due to steady-state surface contamination at the 1×10^{-4} Torr working pressure.

One of the most surprising results from the single-chamber apparatus is that the SEELEM spectra recorded on Au (5 eV work function) and Cs (2 eV work function) are essentially identical. The simplest explanation of this result, which is independently confirmed by statistical measures applied to simultaneously recorded SEELEM and UV-LIF spectra, is that both Au and Cs surfaces are vastly more sensitive to S_1 than triplet character. Experiments carried out during 2001 show that adsorbed acetylene shuts down triplet detectivity.

The second-generation apparatus consists of two (source and detector) independently pumped chambers, a new pulsed nozzle with a 35 μ s FWHM open time, a skimmer located ~ 1 cm (adjustable) downstream from the nozzle, the distance from excitation region to SEELEM surface variable from 3 to 48 cm (plus an additional 33 cm with addition of an auxiliary chamber), beam-on pressure in the detector chamber of 4×10^{-6} Torr (in contrast to 1×10^{-4} Torr with the nozzle operated at 10 Hz in the original apparatus), optimized f/2 optics for both IR-LIF and UV-LIF, and an IR detector (same D^* as the previous IR detector) with a response time (8 μ s now vs. 100 μ s previously) sufficiently fast to measure the radiative lifetime of the unknown carrier of the IR-LIF signal (most likely the \tilde{A} -state of CCH). The improved collection optics and IR detector response should increase the IR-LIF signal by a factor of ~ 50 . The use of a skimmed jet will reduce the Doppler widths of the IR-LIF and UV-LIF channels to ~ 0.003 cm^{-1} , which in turn motivates replacement of our pulsed dye laser by an available pulse-amplified, cw tunable laser system.

2. The Mechanisms of the SEELEM and UV-LIF Detection Processes.

The SEELEM signal strength is given by the product of three factors: excitation probability, probability that the molecular excitation survives through the transit time from laser excitation to impact on the SEELEM surface, and probability of ejection of an electron from the detector surface [HUM97]. The SEELEM signal that results from excitation of the j -th eigenstate will depend on the specific admixture of electronic-vibrational basis states in that eigenstate:

$$|j\rangle = \alpha|S_1, v\rangle + \sum_i \beta_i|T_3, v_i\rangle + \sum_i \gamma_i|T_1, v_i\rangle + \sum_i \delta_i|S_0, v_i\rangle \quad (1)$$

where α , β , γ , and δ are mixing coefficients which express the respective contributions of S_1 , T_3 , T_1 , and S_0 electronic states to the mixed eigenstate and the subscript i labels the vibrational level of the corresponding electronic state. All of the mixing coefficients are real, and

$$1 = |\alpha|^2 + \sum_i \left[|\beta_i|^2 + |\gamma_i|^2 + |\delta_i|^2 \right] \quad (2)$$

Since the electronic ground state is a singlet, S_0 (total electron spin $S = 0$), and electronic transitions between singlet and triplet (total electron spin $S = 1$) states are forbidden, only the $S_1 \leftarrow S_0$ transition is allowed and the S_1 basis state is called the “*bright state*.” All of the other basis states, because they can neither be excited from S_0 nor decay into S_0 by spontaneous fluorescence, are called “*dark states*”. The T_3 state is a special triplet state because it is a more highly excited electronic state than T_1 . Because T_3 is more highly excited than T_1 , the density of vibrational levels associated with the T_3 electronic state (~ 1 per 100 cm^{-1}) is much smaller than that associated with T_1 (10 per cm^{-1}) at the energy of the initially prepared S_1 *bright state*. Therefore, it is common to consider only one T_3 vibrational level as contributing significantly to the j -th eigenstate, and this unique T_3 vibrational level *might* have special dynamical significance. It is called the “*Doorway State*” because it *could* play a promoter role in the coupling between S_1 and the dense manifold of dark states, T_1 and S_0 . Therefore, in what follows, mixing coefficients for only one S_1 (α) and one T_3 (β) basis state will be considered in Eqs. (1 and 2).

The three multiplicative factors that give the relative intensity of a transition terminating on the j -th eigenstate in a SEELEM spectrum are:

$$\text{Excitation Probability} = I S_{J',J''} N_{J''} \alpha^2 \quad (3a)$$

$$\text{Survival} = e^{-[\alpha^2 \Delta t / 0.27]} \quad (3b)$$

$$\text{Electron Ejection} = \left[\alpha + \theta_3 \beta + \theta_1 \sum_i \gamma_i \right]^2 \quad (3c)$$

where I is the laser intensity, $S_{J',J''}$ is the rotational linestrength (Hönl-London factor), $N_{J''}$ is the population of the lower level of the transition, Δt is the flight time (in μs) from excitation to detector plate, 0.27 μs is the radiative lifetime of a pure S_1 basis state [OCH87, 91, ABE93], and θ_3 and θ_1 are unknown factors that express the *amplitude* detectivities of T_3 and T_1 relative to S_1 . Notice that, although the first and third factors increase as the fractional S_1 character increases, one can have too much of a good thing. The survival factor typically prohibits SEELEM detection of eigenstates with more than $\sim 1.4\%$ S_1 character ($\tau < 2 \times 10^{-5}$ s). Our original expectation was that the relative detectivities of S_1 and T_3 on Au and of S_1 , T_3 , and T_1 on Cs would be nearly identical (θ_3 and $\theta_1 \approx +1$ or -1). This would have greatly facilitated direct experimental measurement of the fractional T_3 and T_1 characters and also have led to a factor of ~ 1000 enhancement of the SEELEM intensity on Cs relative to that on Au! The expected 1000-fold enhancement on Cs reflects the $T_1:T_3$ ratio of vibrational densities of states at the 5.5 eV energy of the S_1 $3v_3$ bright state. However, despite the well documented triplet-dominated sensitivity of SEELEM detectors to eigenstates with fractional singlet character (α^2) less than 0.01% [LIS80, SNE88, 91], the SEELEM spectra recorded in our first-generation apparatus are compatible with *zero* sensitivity to triplet character. There is a fundamental reason why the SEELEM process should be more sensitive to S_1 than triplet character and the sensitivity to triplet character should be more rapidly degraded by contamination [HAG54]. Recent experiments on a clean Cs surface yielded the predicted 1000-fold enhancement over the Au surface.

Two distinct physical processes contribute to the SEELEM effect [HAG54]: (i) an electron is ejected from the metal surface when the molecule transfers its electronic excitation energy to the surface via a long-range interaction between the molecular transition dipole moment and the image dipole moment in the surface [CHA78]; (ii) an electron is ejected from the molecule when, via orbital overlap at short-range, an electron is transferred from the metal surface into a vacancy in the lowest lying unfilled molecular orbital. The first process is exclusively sensitive to S_1 character and relatively robust with respect to surface contamination; the second is comparably sensitive to S_1 and T characters but rapidly

degraded by molecules adsorbed on the detector surface. Both processes require that the vertical electronic de-excitation energy in the molecule is larger than the work function of the metal.

The product of three factors also gives the relative intensities of transitions in the UV-LIF spectrum:

$$\text{Excitation Probability} = I S_{J',J''} N_{J''} \alpha^2 \quad (4a)$$

$$\text{Non - flyout} = (1 - e^{-\alpha^2 \delta t / 0.27}) \quad (4b)$$

$$\text{PMT Cathode Efficiency} = f(\lambda) \quad (4c)$$

where the excitation probability factor is identical to that for the SEELEM spectrum, δt is the time required for the excited molecule to travel out of the PMT field of view, and $f(\lambda)$ is the wavelength-dependent PMT cathode sensitivity. The survival factor in the SEELEM expression (Eq. 3b) and the Non-flyout factor in the UV-LIF expression (Eq. 4b) result in a near complementarity of the SEELEM and UV-LIF channels; eigenstates that are SEELEM detectable ($\tau > 20 \mu\text{s}$, $\alpha < 0.12$) are not UV-LIF detectable and *vice versa* ($\tau < 3 \mu\text{s}$, $\alpha > 0.30$).

Previous studies of the SEELEM effect (often called the "Auger effect" [HAG54, LIS80], but this name has acquired a completely different connotation in Surface Science) in atoms and diatomic molecules have been unable to explore the relative strengths of and quantum interferences between the contributions from multiple electronic states to SEELEM signals [HAG54, LIS80, SNE86ab, 88, 91]. Acetylene is unique in the possibility that two (S_1 and T_3) electronic states potentially contribute to the SEELEM effect on Au and three (S_1 , T_3 , and T_1) on Cs. Acetylene is also unique in that its vibronic density of states is sufficiently low (neglecting interaction with S_0) that individual eigenstates can be resolved in both SEELEM and LIF spectra, and that a complete model for the interactions among these states can be inferred from the totality of spectroscopic observations [DRA94, OCH87, 91, DUP91, 93, 95ab, HUM97]. Our multispectral studies of acetylene will determine the signs and magnitudes of the amplitude detectivity factors in Eq. 3c, thereby providing fundamental insights into the mechanism of the SEELEM effect and the factors that determine the quantum efficiency of SEELEM detection on various metal surfaces. In turn, a fundamental understanding of the SEELEM effect, in particular of the interference effects among several electronic states in a mixed eigenstate, in combination with the high resolution multispectral data sets recorded in our second-generation apparatus, will yield qualitatively new insights into the mechanisms of Intersystem Crossing and Internal Conversion in small polyatomic molecules [BIX67, AVO77, STA81, NES96].


3. The Unexpectedly Sensitive Yet General Near Infrared Fluorescence Detection Channel.

We first observed near infrared fluorescence resulting from UV laser excitation of acetylene in static gas cell experiments [DRU97]. Careful examination of the laser intensity dependence of the IR signal suggests that most of the IR fluorescence is due to a multiphoton process that produces an unknown electronically excited photofragment of acetylene. This multiphoton process is enhanced at the one-photon level by resonance with an eigenstate of predominant S_1 $3v_3$ character. In the static gas cell experiment, the nearby S_1 $4v_b$ perturber levels (four quanta of nontotally symmetric bending vibration [SCH86, UTZ93, CUN00a]) did not contribute significantly to this multiphoton process.


In a serendipitous and improvised (sub-optimal $\sim f/10$ light collection and too slow 100 μs time response of the IR detector) experiment, we discovered that the near-IR-LIF signal in our first-generation molecular beam apparatus is respectively stronger than the SEELEM signal and comparable to the UV-LIF

signal and nearly identical in spectral pattern to that in the UV-LIF channel (Figure 3). This implies that the quantum yield (signal counts per absorbed UV photon) for two-photon excitation of an infrared-emitting photofragment must be at least comparable to that for direct, one-photon excitation of UV-LIF detectable acetylene molecules. The solid angle of the SEELEM detector is such that all molecules in the jet, which pass through the UV-laser excitation region, will strike the active area of the SEELEM detector. However, we have crudely measured the quantum efficiency for SEELEM detection to be $\sim 5 \times 10^{-6}$, much smaller than the quantum efficiency of the infrared detector. The product of detector solid angle and efficiency factors for SEELEM vs. IR-LIF detection favors IR-LIF by a factor of ~ 100 , which implies that the strengths of the two excitation pathways will be comparable at UV laser pulse intensities only slightly larger than the unfocussed 100 μJ currently in use. We believe that this is not an accident and that the resonance enhanced, multiphoton IR-LIF channel will be an unexpectedly sensitive detection channel for characterizing metastable states and Intersystem Crossing in many small polyatomic molecules which have a multiple bond.

Figure 3. Comparison of UV-LIF (upward) and IR-LIF (downward) spectra in the region of the S_1 $3\nu_3$ level.



In acetylene, a single $\pi^* \leftarrow \pi$ orbital excitation produces the S_1 and $T_{1,2,3}$ excited states [ING53, INN54, PRI45, KAM70, DEM75, WET78, SEK85, LIS86, VAC93, YAM93, 94, STA94, SHE96, CUI96]. A second orbital excitation would nominally lead to a CCH photofragment with a hole in the π orbitals or a singly occupied π^* orbital or both. These correspond to electronically excited states of CCH. If the excited state is a doublet state ($S=0.5$), then the excited state could decay by near infrared fluorescence analogous to the $\text{CN } A^2\Pi \rightarrow X^2\Sigma^+$ transition. If the excited state is a quartet state ($S=1.5$), it would be sufficiently long-lived and energetic to be SEELEM detectable. The metastable photofragment that was observed in the TOF-SEELEM spectrum in the region of S_1 $4\nu_3$ [HUM97] *could* be a quartet state of CCH.



It is reasonable to expect, for a large and important class of unsaturated polyatomic molecules, that sequential excitation processes, via *either the low- ν S_1 or low- ν T character* of one-photon excited eigenstates, will *generally* give rise to a near IR fluorescence signal that is of comparable intensity to the SEELEM signal. It is important to note that the information in the SEELEM and IR-LIF channels is complementary. SEELEM detectivity should be independent of T vibrational state (neglecting mode-specific fractionation of T into S_0), whereas the IR-LIF channel depends on Franck-Condon overlap and energy match with a predissociated, gerade-Rydberg vibronic final state at the two-photon level [ASH87, TAK92]. S_1 and T basis states of low vibrational excitation (in which one turning point resembles the equilibrium geometry of the final dissociative state) should dominate the IR-LIF signal channel, provided that the UV photon is resonant with the g-Rydberg $\leftarrow (T, S_1)$ transition. In contrast, of those molecules that absorb one photon, only a small fraction will escape all of the fates antithetical to SEELEM detection: absorption of a second photon followed by rapid radiative decay or dissociation into electronically unexcited photofragments, rapid UV fluorescence decay of the one-photon excited eigenstate prior to impact on the SEELEM detector (too much S_1 character), and failure of the one-photon excited eigenstate to eject an electron from the SEELEM detector (too little S_1 character).

IR-LIF spectra were recorded in the region of the HCCH S_1 zero-point level, which is also the predicted energy region [CUI96, SHE96, VAC93, 96] of the minimum energy half-linear point on the T_2 potential energy surface. Three vibrational bands, the acetylene $S_1 \leftarrow S_0$ origin band and two $S_1(1\nu_3) \leftarrow S_0(2\nu_4)$ hot bands, are observed in both UV-LIF and IR-LIF channels, but the *hot bands* were ~50 times *stronger* in the IR-LIF than UV-LIF channel. This observation is the basis for our claim that Franck-Condon and energy match factors determine IR-LIF intensities.

Calculations [CUI96] suggest that $T_2 \sim T_1$ spin-orbit interactions are much weaker than the $T_3 \sim T_2$ and $T_3 \sim T_1$ interactions. This is supported by Kanamori's [KAN98] observation of negligible fractionation of the upper level of the $T_2(cis) \leftarrow T_1(cis)$ origin band. Therefore we expect that, below the energy of the T_3 zero-point level, the T_2 vibrational levels will be sufficiently free of fractionation into high vibrational levels of S_0 that several vibrational levels of the T_2 state (especially overtones of the *trans*-bend in the *trans*-bent isomer and overtones of the *cis*-bend in the *cis*-bent isomer) should be observable in the IR-LIF spectrum. On the other hand, we do not expect to see transitions into the denser manifold of T_1 vibrational levels in the S_1 origin band region of the IR-LIF spectrum, because those states are too highly excited (and

extensively mixed by anharmonic resonances) to have appreciable Franck-Condon overlap with the final (linear geometry) state that dissociates into the IR-emitting photofragment. Our IR detector failed just after our initial unsuccessful attempt to record $T_2 \leftarrow S_0$ IR-LIF spectra. The first-generation apparatus was dismantled and the second-generation molecular beam apparatus was designed to optimize the IR-LIF detection channel.

$T \leftarrow S_0$ transitions in H, C, N, O molecules are extremely weak. Previous efforts to detect such transitions have focussed on detection of extremely weak phosphorescence (direct or sensitized [SUZ97, PRA88]) or SEELEM signals from metastable parent molecules. Multiphoton photofragmentation processes will frustrate attempts to enhance the intensity of such parent-metastable-based signals by increasing laser intensity. Since the transition moment for the $T \leftarrow S_0$ transition will be enormously smaller than those for all subsequent $T' \leftarrow T$ transitions, a very small fraction of the molecules excited from S_0 will survive the excitation pulse as intact parent metastable molecules. The IR-LIF detection scheme, which is intrinsically a multiphoton photofragmentation process, will be far more robust and generally applicable than the parent molecule phosphorescence and SEELEM detection schemes!

4. Statistical Measures.

A central goal of this project is to gain information about a large and poorly understood class of long-lived, energetic, and non-fluorescing molecular states that cannot be obtained by known spectroscopic techniques. Such a goal inevitably requires uncovering and dealing with unprecedented spectral complexity. We have built an apparatus with which we have demonstrated the capability to record enormous multispectral TOF-SEELEM/UV-LIF/IR-LIF data sets. It would be naïve to expect that these data could be directly input into a least-squares-fitting routine that is based on a standard spectroscopic effective Hamiltonian model [LEF86]. Moreover, standard spectroscopic models are likely to be prohibitively opaque when applied to energy regions where the vibrational density of states approaches 10 per cm^{-1} .

We have devised several kinds of statistical pattern recognition schemes to deal with the unique problems posed by the data sets we are now capable of recording [COY95, 97, JAC97, ALT99, 00ab, CUN00b].

Whenever a dynamical quantity is conserved (either rigorously, as for a symmetry-based good quantum number, or approximately, as defined by the finite time that corresponds to the resolution of the spectrum), that conserved quantity will give rise to a pattern that is exactly repeated in the spectrum. The difficulty that must be overcome is that the patterns associated with different values of the conserved quantity (e.g. the rotational quantum number, J) often overlap each other and these overlapping patterns must be disentangled, without benefit of any prior knowledge about the nature of the patterns. We have developed powerful spectral autocorrelation and cross-correlation methods [JAC97, COY97, CUN00b] and applied them to a variety of problems [JAC98, VIT97a-g].




Figure 5. Intensity weighted average energies, extracted by the XAC pattern recognition scheme, from extensively overlapped fractionated J' -patterns in UV-LIF and SEELEM spectra in the region of S_1 $3\nu_3$. With the exception of the $J'=4$ point, the J' -sorting procedure is perfect, and the UV-LIF spectrum yields the horizontal straight line required by perturbation theory, with scatter on the order of the 0.08 cm^{-1} resolution. The larger deviations, at $J=2$ and 3 in the SEELEM spectrum, are genuine and constitute proof that the $S \sim T$ coupling is *Doorway-Mediated* rather than *Direct* and that the T_3 doorway state overtakes S_1 from below at $J=3$.

Figure 5 displays intensity weighted average rotational energy plots obtained from the UV-LIF and SEELEM spectra of the acetylene $\tilde{A} - \tilde{X} \ V_1^3 \ K_1^0$ hot band by the eXtended spectral AutoCorrelation (XAC) method. The sorting of the spectra into separated patterns associated with each of the upper state J' -values is satisfactory, but would be improved if the spectra could be recorded at higher resolution and signal:noise ratio. Significantly, we observed that an improvement in UV laser resolution (0.2 cm^{-1} at UCSB [HUM97], 0.08 cm^{-1} at MIT [ALT98, CUN00b]) led to a dramatic S:N improvement. We plan to increase the laser resolution from the present $\sim 0.08 \text{ cm}^{-1}$ FWHM Nd:YAG pumped, frequency doubled, pulsed dye M to $\sim 0.003 \text{ cm}^{-1}$ by replacing our laser by a frequency doubled, pulse-amplified, cw-doubled, Ti:sapphire laser (Coherent 899-29, supplied by DURIP Grant F499620-97-1-0245) during the requested grant period.

Another kind of pattern serves as the unique signature of two qualitatively opposite mechanisms of Intersystem Crossing: the "*Doorway Mediated*" and "*Direct*" models [AVO77, ALT00ab]. In the *Doorway* model, there is one *bright* state (S_1), one *doorway* state (T_3), and a dense manifold of *dark* states (T_1, S_0). The *bright* state is bright because it is connected to thermally populated levels of the ground electronic state by an electric dipole allowed transition. The *doorway* and *dark* states are dark because they can neither be excited from the ground state nor decay by emission of detectable fluorescence. The *doorway* state is special, acting as a promoter of Intersystem Crossing because it is coupled by off-diagonal spin-orbit matrix elements to *both* the *bright* and *dark* states. In the *Doorway* model the *bright* state has no direct off-diagonal matrix elements to the *dark* states. In the *Direct* model, there is no special promoter or *doorway* state; the *bright* state has nonzero off-diagonal matrix elements to all *dark* states.

Since the statistical measures used to distinguish qualitatively between the *Doorway Mediated* and *Direct* models also serve to determine quantitatively the molecular constants of the *doorway* state and the *bright-doorway* state coupling matrix element [ALT00ab], both the qualitative and quantitative capabilities of each statistical measure are discussed here. The statistical measures are:

(i) Mean Energy.
$$E_{ave} = \sum I_i E_i \quad (5)$$


(ii) Skewness
$$S = \sum_i I_i^2 [E_i - E_{ave}] \quad (6)$$

(iii) Correlation.
$$C_{\Delta\nu} = \frac{\sum_i [I_i^{LIF} * I_i^{LEM}]_{\Delta\nu}}{\sum_i [I_i^{LIF}]^2} \quad (7)$$

(iv) Fractionation.
$$F = \sum_i I_i^2 \quad (8)$$

(v) Lawrance-Knight De-Diagonalization [BER76, LAW85, 91, LEH91].

I_i is the intensity in the i -th resolution element of a spectrum, E_i is the energy that corresponds to the i -th resolution element, E_{ave} is the intensity weighted energy averaged over all resolution elements in a spectrum (e.g. SEELEM, UV-LIF, or IR-LIF) which belong to the same value of the rigorous quantum numbers (e.g. J' and parity), and $[I_i^{LIF} * I_i^{LEM}]_{\Delta\nu}$ is the product of intensities in LIF and SEELEM spectra calculated at resolution $\Delta\nu$.



A good framework for understanding the information revealed by each of the statistical measures is to imagine, for the *Doorway Mediated* model, that the *bright* and *doorway* basis states have slightly different rotational constants and vibrational energies such that, on an energy vs. $J(J+1)$ plot, the straight line of slope B_{bright} intersects the straight line of slope $B_{doorway}$ at J_c . At the level crossing, J_c , the resultant eigenstates exhibit the largest level-shifts from the unperturbed basis state energies and also the largest *bright-doorway* mixing coefficients. Also, a sign-reversal of mixing coefficients and quantum interference effects occurs on opposite sides of J_c [LEF86]. If the coupling is *Direct* rather than *Doorway Mediated*, then there will be no sign of a level crossing nor, for any value of J' , any sign of asymmetries in the various spectra associated with a *doorway* state either above or below the *bright* state.

(i) Mean Energy. In the UV-LIF spectrum, the J' -dependence of E_{ave} is indistinguishable from the energy of the *bright* state for both *Direct* and *Doorway* models [ALT00a]. This is useful because it provides a simple and accurate method for extracting the molecular constants of the *bright* state from a complicated UV-LIF spectrum provided that the J' -sorting procedure is successful. In a *Doorway* model SEELEM spectrum, the J' -dependence of E_{ave} exhibits a strong asymmetry, which locates J_c and for each- J' determines whether the *doorway* state lies above or below the *bright* state. In a *Direct* model SEELEM spectrum, there is an approximately constant offset of E_{ave} from the energy of the bright state.

(ii) Skewness. Only the *Doorway* model SEELEM spectrum exhibits skewness [ALT00a]. The skewness is zero at J_c and the sign of the skewness determines whether the *bright* state lies above or below the *doorway* state.

(iii) Correlation. The correlation depends strongly on Δv , approaching zero as Δv goes to zero. For the *Doorway* model, at finite resolution, the correlation increases from nearly zero at J_c to near unity at J' values above and below J_c . For the *Direct* model, the correlation is small and essentially independent of J' [ALT00a].

(iv) Fractionation [HEL97, NES96]. The fractionation index can range from 1 (no fractionation) to $1/N$ (maximum fractionation), where N is the number of potentially interacting states. The *Doorway* model exhibits a characteristic non-symmetric fractionation pattern with minimum fractionation index (approximately equal to the roughly constant value for the *Direct* model) near J_c in both UV-LIF and SEELEM spectra. The absence of reflection symmetry about J_c is due to quantum interference between S_1 amplitudes obtained directly and via the T_3 (*doorway*) perturber [ALT00a].




Figure 7. Summary of four statistical measures applied to *Doorway Mediated* and *Direct Model* spectra [ALT00a]. The patterns shown here illustrate the qualitative differences between *Doorway Mediated* and *Direct Model* behavior and also show how quantitative molecular constants for the *Doorway* state may be obtained without a full fit of eigenstates to an H^{eff} model.

(v) Lawrance-Knight De-Diagonalization. The Lawrance-Knight (L-K) method, originally proposed by Berg [BER76], later reformulated by Lawrance and Knight [LAW85, 91], and critiqued by Lehmann [LEH91], permits exact inversion of a spectrum (frequencies and relative intensities) to a special

kind of zero-order model (one bright state coupled to a “prediagonalized” manifold of dark states), which is identical to our *Direct* model. One obtains the deperturbed energies of the *bright* and *dark* states and the values of the coupling matrix elements between the *bright* state and each *dark* state. When the L-K method is intentionally applied to the *Doorway* model case (for which it was explicitly not intended), one obtains information (from the skewness of the L-K output) about the location of the *bright* state ~ *doorway* state level crossing, J_C , whether the *bright* state is crossed by the *doorway* state from above ($B_{\text{bright}} > B_{\text{doorway}}$) or below, and the magnitude of the *bright* ~ *dark* coupling matrix element [ALT00b].

The L-K procedure is the most powerful of all the statistical measures we have examined, both for qualitatively distinguishing between *Doorway* and *Direct* model coupling cases and for quantitatively determining the molecular constants that describe the *doorway* state and the *bright* ~ *dark* coupling matrix element. Its value as a spectroscopic tool is illustrated by the following example where we were forced to revise our original model for the SEELEM detection process [ALT00b].

Equation (3c) gives the probability of ejection of an electron from the metal surface of the SEELEM detector. The form of this expression, where an event probability is given by the square of a sum of amplitudes, is demanded by fundamental postulates of Quantum Mechanics. However, the signs and magnitudes of the θ_1 and θ_3 factors are dependent in an unknown way on the electronic wavefunctions of the various molecular electronic states and the metal surface. We assumed initially that both θ factors are approximately +1 or -1. When we make this assumption, several inconsistent conclusions are drawn from UV-LIF and SEELEM spectra. The L-K skewness measure, applied to the high resolution UV-LIF spectrum recorded by Drabells et al. [DRA94] implies that the T_3 *doorway* state starts out below the S_1 $3v_3$ $K'=1$ *bright* state at $J'=1$ and crosses the *bright* state at $J_C=3$. The skewness statistic defined by Eq. 6, applied to the S_1 $3v_3$ $K'=1$ *bright* state SEELEM spectrum, implies that the *doorway* state starts out above the *bright* state at $J'=1$. However, when θ_3 is reduced from 1 to 0.01, the sense of the SEELEM skewness is reversed and the statistical analyses of the SEELEM and UV-LIF spectra give consistent results [ALT99, 00b]. A much reduced detectivity factor for triplets relative to S_1 also explains the absence of any significant differences between both the shapes and the relative intensities of the SEELEM spectra recorded on Au (where T_3 is in principle detectable) and Cs (where both T_3 and T_1 are in principle detectable). It is plausible that the Au/Cs invariance of the SEELEM spectra is due to contamination of the Cs surface, because SEELEM detection of triplet states requires a shorter-range interaction between molecule and metal surface than does detection of singlet states [CHA78].


5. Dispersed Fluorescence Studies of Spectroscopic Perturbations in HCCH S_1 $3v_3$.

Spectroscopic perturbations are windows through which one can view otherwise unobservable classes of electronic-vibration states [LEF86] and, most significantly yet least appreciated, the *causal mechanisms* (in contrast to phenomenological rates) of dynamical processes such as Intramolecular Vibrational Redistribution (IVR), Intersystem Crossing (ISC), and Internal Conversion (IC) [BIX67, AVO77, STA81, NES96]. Rather than being sought out as sources of unique information, perturbations are generally given a wide berth by experimentalists. The reason for this is simple: a perturbation destroys the regular pattern of transition frequencies and intensities on which the assignment of a spectrum is based [LEF86]. A central theme of this project is the development of multispectral techniques and pattern-recognition methods that will enable experimentalists to record and interpret dynamically relevant spectra, regardless of the apparent complexity of these spectra.

A perturbation disrupts the pattern in an absorption or LIF spectrum. Dispersed Fluorescence (DF) spectroscopy, wherein one records the spectrum of the fluorescence that results when a single eigenstate in an LIF spectrum is excited, contains information that can reveal the identity of several of the electronic-vibration basis states present in the selected eigenstate [YAM91, SOL95, 96, ACE99, CUN00a].

There is a long history of observations and partial assignments of the low-J perturbers of the HCCH S_1 $3v_3$ $K=1$ manifold [SCH86, UTZ93, DUP91, 93, 95b, OCH87, 91, WOL90, DRA94]. It is clear that at least five classes of basis state are involved: (i) the S_1 $3v_3$ (v_3 is the *trans-bend*) LIF-bright state, (ii) the S_1 $4v_b$ class of LIF-dark but DF-bright anharmonic ($4v_4$, $2v_4 + 2v_6$, and $4v_6$, where v_4 is the torsion and v_6 is the antisymmetric in-plane bend) and Coriolis ($v_4 + 3v_6$ and $3v_4 + v_6$) perturbors [UTZ93], (iii) the T_3 *doorway* state, which is both LIF- and DF-dark [CUN00a], (iv) the T_1 dark states [DRA94], and (v) high vibrational levels of S_0 [DUP91, 93, 95b]. The DF spectrum from an unperturbed S_1 nv_3 level is well known, from our DOE-supported project [YAM91, SOL95, 96, ACE99], to consist of a progression of *polyads*, each illuminated by a single *bright-state* (which involves excitation exclusively in the CC-stretch and *trans-bend* vibrations of S_0). The two distinct patterns of frequencies and relative intensities within each polyad, generated by *either* a $3v_3$ or a $4v_b$ pluck, are known from the H^{eff} model that provides a complete picture of the IVR on the acetylene S_0 potential surface [JAC99, CUN00a]. When the eigenstate from which the DF spectrum is recorded contains significant contributions from both $3v_3$ and $4v_b$ basis states, the observed intrapolyad pattern is a superposition of the $3v_3$ and $4v_b$ patterns. However, when the selected eigenstate contains T_3 , T_1 , or S_0 basis state character, the result is no distortion of either the intrapolyad or interpolyad patterns in the DF spectrum. The intensity pattern in our supersonic jet UV-LIF spectrum implies that the S_1 $3v_3$ bright state primarily interacts with two ($J = 1, 2$, and 3) or three ($J = 4$) dark states [CUN00a], Drabbels' higher resolution UV-LIF spectra [DRA94] show that each of these is fractionated into ~ 10 T_1 quasi-eigenstates, and the Zeeman Anticrossing (ZAC [DUP91]) and Quantum Beat (ZQB [DUP93, 95b]) spectra show that each quasi-eigenstate is capable of interacting with as many as 100 dark S_0 states. Our DF spectra show that the S_1 $3v_3$ bright state and the T_3 *doorway* state cross at $J_c = 3$ and that the level crossing between $3v_3$ and $4v_b$ occurs at $J_c < 1$. Rotational constants and perturbation matrix elements are also determined [ALT00b, CUN00a].

Figure 8. The S_1 $3v_3$ bright state is perturbed by T_3 and by S_1 $4v_b$ and at least one other unidentified state. The labels T, M, and E refer respectively to triplet (T_3), main (S_1 $3v_3$), and extra (S_1 $4v_b$).




The DF results [CUN00a] confirm our statistical analyses of our SEELEM and UV-LIF spectra [ALT00a] and our L-K analysis of Drabbels' high-resolution UV LIF spectra [ALT00b]. They also explain the plateau intensity pattern observed in our UCSB SEELEM spectra [HUM97]. The plateau is the result of a quantum interference effect. The interference switches from constructive to destructive at $J=3$ because of a sign-reversal of the $S_1 \sim T_3$ mixing coefficients at J_c . The eigenvectors of our $T_3 \sim 3v_3 \sim 4v_b$ 3-state fit

model confirm an essential qualitative feature of our ISC model. The T_3 *doorway* state interacts primarily with the S_1 $3v_3$ *bright* state but is prohibited, by poor Franck-Condon overlap, from interacting with the S_1 $4v_b$ perturber. Therefore the fractionation (into the ~ 10 state per cm^{-1} manifold of T_1 *dark* states) of each feature in the UV-LIF spectrum [DRA94] correlates with the fractional T_3 *doorway* state character in that feature: the nominal T_3 feature is most fractionated and the nominal S_1 perturber feature is least fractionated. This is *mode-specific* ISC!

This is mechanism, not phenomenology. There is a Franck-Condon reason why a special class of *doorway* state is effective in initiating the nonradiative decay of each class of *bright* state.

6. UV Cavity Ringdown.

We plan to exploit the exceptional sensitivity and simplicity of Cavity Ring-Down Spectroscopy (CRDS) [OKE95, ZAL95, LEH96, HAG98] for direct observation of the $T_1 \leftarrow S_0$ absorption spectrum. However, in the $\lambda < 300$ nm UV region, mirrors with reflectivity greater than 99.8% are either unavailable or prohibitively expensive. In a DOE-supported project we have implemented a CRDS scheme that closely approaches the quantum noise limit ($N^{0.5}$ where N is the number of photons/pulse incident on the entrance mirror of the cavity) even when 99.5% reflective mirrors are used. The critical features of this scheme are (i) the length of the CRD cavity is chosen so that the output beams from the cavity sequentially illuminate three stationary spatial regions on the photodetector surface and (ii) our least squares fitting algorithm takes into account the existence of three families of exponentially decaying outputs that have the same decay rate but different amplitudes.



Ringdown signal recorded for "tuned" cavity with three "families" of peaks.

Figure 9. UV Cavity Ringdown. Owing to the different sensitivity of the three spots on the detector illuminated by the output beams from the ringdown cavity, there are three families of decay curves, each with the same exponential decay constant. The pulses inside the cavity are resolved because the 12 ns round trip time inside the 2m cavity is longer than the ~ 4 ns pulse.

inside the cavity are resolved because the 12 ns round trip time inside the 2m cavity is longer than the ~4 ns pulse.

A crude estimate of the intensity of a direct $T_i \leftarrow S_0$ absorption band is obtained from that of the corresponding $S_1 \leftarrow S_0$ band [ING53, NAK64], multiplied by the fractional S_1 character in T_i obtained using nondegenerate perturbation theory for the first-order correction to the wavefunction [LEF86]. The spin-orbit matrix element is taken as the spin-orbit coupling constant for the Carbon 2p orbital, $\zeta = 29 \text{ cm}^{-1}$ and the energy difference between S_1 and T_i states is taken as $10,000 \text{ cm}^{-1}$. Thus the fractional S_1 character in T_i will be 8×10^{-6} and the strongest $T_i \leftarrow S_0$ absorption band will therefore be on the order of 10^5 times weaker than the strongest $S_1 \leftarrow S_0$ band. As practise for our first attempt at recording a $T_1 \leftarrow S_0$ band by CRDS, we recorded the $S_1 \leftarrow S_0$ origin band at $\lambda = 236 \text{ nm}$, which has a vibrational band oscillator strength of $f = 2.5 \times 10^{-9}$, a factor of ~700 smaller than that of the V^3_0 (" $3v_3$ ")band [ING53, WAT95].

It will be interesting to compare the relative sensitivities and unique electronic-vibration basis state selectivities of IR-LIF, SEELEM, and CRDS for recording direct $T \leftarrow S_0$ transitions.

7. Hg* Photosensitization Experiments.

There is a long history in Organic Photochemistry [OKA78] of using Hg* photosensitization [HUN71, BUR72, WEN79] to promote anti-Woodward-Hoffmann [WOO71] reactions. 253.7 nm radiation from a mercury lamp is used to excite Hg atoms from the 1S_0 electronic ground state to the 3P_1 excited state that in turn is efficiently transferred into the long-lived 3P_0 state by collisions with a nonreactive gas such as N_2 . The photosensitized reaction is carried out in a quartz tube that contains a trace amount of Hg (1 mTorr), a high pressure of N_2 (1 atm), and the organic reagents (~10 Torr). Efficiencies near 100% for transferring the excitation from Hg (3P_1) to Hg (3P_0) to the unsaturated organic reagent are typically claimed [WEN79].

Preliminary gas cell experiments demonstrate the feasibility of a lower pressure implementation of the Hg* photosensitization scheme. Direct laser excitation in the blue wing of the Hg $^3P_2 \leftarrow ^1S_0$ or $^3P_0 \leftarrow ^1S_0$ atomic transition, in the presence of 100 Torr of Xe, results in a large and spatially uniform population of long lived (~1 ms) 3P_2 or 3P_0 Hg atoms, respectively [OHM94, 95, 99, KUR98]. The populations in the $J = 0, 1$, or 2 fine structure levels of the Hg 3P_J state are monitored by laser excitation of the $^3S_1 \leftarrow ^3P_J$ atomic transition, which results in visible $^3S \rightarrow ^3P$ and UV $^3P_1 \rightarrow ^1S$ fluorescence.

Our plan is to use Hg* photosensitization to populate metastable states of several small molecules, starting with NO and acetylene. The goal is to develop an Intracavity Laser Absorption Spectroscopy (ICLAS) scheme for monitoring populations in metastable electronically excited states of molecules [ICL00]. ICLAS is a versatile, absorption-based scheme, with sensitivity comparable to CRDS, but with two crucial advantages over CRDS: it is a multiplex scheme, capable of simultaneously recording several tens of cm^{-1} of spectrum at 0.02 cm^{-1} resolution, and, in a double-sampling scheme recently implemented by Kachanov in the Steinfeld lab at MIT, ICLAS can be used to monitor kinetic processes at a few microseconds time resolution. The MIT ICLAS setup, the result of a DURIP grant to Steinfeld and Field, is based on a Ti:sapphire intracavity gain medium. The lasing range of Ti:sapphire, 700-1100 nm, is a nearly perfect match for a class of electronic transitions that is ideally suited for monitoring the populations in metastable electronic states. For almost all unsaturated H, C, N, O molecules, the $\pi^* \leftarrow \pi$ or $\pi^* \leftarrow n$ metastable excited state can be monitored via a $\pi \leftarrow \sigma$ or $n \leftarrow \sigma$ electronic transition in the red to near-IR region.

8. Collaborations at AFRL, Hanscom Research Site: Pattern Recognition and Baseline Stripping.

As a result of reciprocal visits by Steven Lipson to MIT and Robert Field and Matt Jacobson to the AFRL Hanscom Research Site, several collaborative projects involving pattern-recognition techniques were initiated. Initial results have been reported in two papers [JAC98, RUC00] and seven conference abstracts [VIT97a-g].

In the first paper [JAC98], Hybrid Linear Pattern Analysis (HLP) was used to extract information from atmospheric emission simulation experiments conducted in the LABCEDE facility of the Hanscom Research Site. HLP combines model-based and model-free pattern-recognition techniques. A pulsed electron beam excites CO in a cryogenic chamber and time resolved IR spectra are recorded. The spectra contain information about the vibrational populations and collisional deactivation rates in CO X $^1\Sigma^+$ $v=0-12$. Severe self-absorption in the $1 \rightarrow 0$ band prohibits the use of a model-based least-squares fit to the time-resolved spectra. The initially unknown pattern associated with the $1 \rightarrow 0$ band is recovered by the extended spectral cross-correlation method (XCC) [JAC97, COY97] by assuming that the spectral profile of this pattern does not change with time. All of the other vibrational bands are free of distortion by self-absorption, thus their band profiles can be modeled by standard least squares fitting techniques. HLP recovers the time-dependent populations in all vibrational levels $v = 1-12$ and the time varying spectral profile of the $1 \rightarrow 0$ band in a single, statistically rigorous fit [JAC98]. This paper was named the AFRL Space Vehicles Directorate "Team Publication of the Year" for 2000.

The second paper [RUC00] addresses the classic problem of separating baseline from signal. This becomes an even more important problem when one begins to apply model-free pattern-recognition techniques (e.g. XCC and XAC) to a data stream where the nature and number of patterns to be extracted are initially unknown. Since the model-free techniques begin with a *recursion map* [JAC97, COY97], which is a plot of the intensities in one data record vs. those in another, it is essential to partition the measured intensities into spurious baseline and real signal contributions. Baseline corrupts patterns and vitiates the effectiveness of any pattern-recognition procedure. This collaboration, between an Australian statistician Andreas Ruckstuhl, James Dodd from AFRL, and the MIT group, materialized at a mini-symposium organized by the P. I. on Modern Spectroscopic Techniques at the 1998 American Chemical Society National Meeting in Boston.

A technique entitled Robust Baseline Extraction (RBE) is introduced, which uses techniques of robust local regression to estimate baselines in spectra that consist of sharp features superimposed upon a continuous, slowly varying baseline [RUC00]. The technique is applied to synthetic spectra, to evaluate its capabilities, and to laser-induced fluorescence spectra of OH (produced from the reaction of ozone with hydrogen atoms). The latter example is a particularly challenging case for baseline estimation because the experimental noise varies as a function of laser frequency. RBE enabled Dodd et al [DOD00] at the AFRL Hanscom Research Site to obtain rapid-scan records of OH LIF data by scanning slowly near lines and fast between lines.

Similar pattern-recognition methods were developed by other members of the Lipson group and applied to hyper-spectral (one spectrum for each of an enormous number of spatial locations) problems, such as target recognition and the IR spectra of the Earth's atmosphere recorded by the CIRRUS 1A instrument aboard the Space Shuttle [VIT97a-g].

9. The Spectroscopic Signature of Bond-Breaking Isomerization in HCP.

One of the classic problems in spectroscopy and molecular dynamics is: “what happens to the quantum spectrum and to classical trajectories when bond-breaking isomerization becomes energetically accessible?” The naïve expectation was that spectra would become uninterpretable complex and that trajectories would become chaotic. Even though HCP corresponds to a saddle-point rather than a local minimum on the HCP S_0 potential energy surface (PES), the HCP \leftrightarrow HPC system has emerged as the simplest and best understood (experimentally and theoretically) bond-breaking isomerizing system [HCP99].

The HCP project [LEH85] began at Princeton and was joined at MIT in 1989 as an AFOSR-supported Simulated Emission Pumping (SEP) study of vibrationally highly excited levels of the HCP $X \sim {}^1\Sigma^+$ electronic ground state [CHE90a]. A postdoc, Dr. Haruki Ishikawa [ISH96] made critical progress in the interpretation of the SEP spectra. When Dr. Ishikawa returned to Tohoku University, a long term MIT-Tohoku collaboration was established [HCP99]. The experimental part of the HCP project and the fitting of all spectra to Ishikawa’s polyad H^{eff} model were transferred to Prof. Ishikawa’s laboratory. Since then several theorists have joined the collaboration (Reinhard Schinke, Stavros Farantos, Marc Joyeux, Michael Kellman, Alejandro Frank and Renato Lemus). They computed a refined *ab initio* PES, calculated wavefunctions and spectroscopic properties of the eigenstates of the PES, examined the nodal structure of these eigenstates, calculated classical trajectories, and subjected the trajectories to the diagnostics of nonlinear classical dynamics [BEC97].

One of the most remarkable things to emerge is that the standard, spectroscopically determined, H^{eff} matrix model contains all of the qualitative changes in spectral patterns, nodal structures, and classical trajectories [HCP99]. Somehow the H^{eff} “knows” the minimum energy isomerization path on the PES! To a very good approximation, even in the energy region of the top of the isomerization barrier, polyads still exist, even though the clustering of vibrational eigenstates into closely spaced polyad groups of levels no longer exists. At high energy the polyads begin to overlap each other, but the inter-polyad matrix elements between the lowest members of a higher energy polyad and the highest members of a lower energy polyad start out very small. This means that the polyad model generates the essential information that ensures its bootstrapping extension into higher energy regions where interpolyad interactions are locally severe but globally modest.

Spectroscopists observe spectra. They cannot directly view wavefunctions, thus their spectroscopic assignments cannot be guided by the nodal structures of observed eigenstates. Their assignments are based on patterns of energy levels and relative intensities, and these assigned levels are the bricks and mortar from which an H^{eff} matrix model is built. This model is a better tool, than *anyone* would have imagined, for predicting and describing unimolecular dynamics, extracting useful information from changes in spectral patterns as the isomerization region is approached, and generating classical trajectories in a *reduced-dimension* space that is much better suited than the exact Hamiltonian to the visualization diagnostics of nonlinear dynamics [HCP99, ACE99].

This study of HCP is the best possible illustration of how the central tool of the molecular spectroscopist, the H^{eff} matrix model, facilitates extraction of **causal mechanisms for dynamics** from spectra of extreme complexity.

D. References

- ABE93. H. Abe and H. Hayashi, "Nonradiative Process Induced by External Magnetic Fields in the C_2H_2 A^1A_u State Near the Predissociation Limit", *Chem. Phys. Lett.* **206**, 337 (1993).
- ACE99. M.P. Jacobson, C. Jung, H.S. Taylor, and R.W. Field, "State-by-State Assignment of the Bending Spectrum of Acetylene at $15,000\text{ cm}^{-1}$: A Case Study of Quantum-Classical Correspondence," *J. Chem. Phys.*; M.P. Jacobson, R.J. Silbey, and R.W. Field, "Local Mode Behavior in the Acetylene Bending System," *J. Chem. Phys.* **110**, 845 - 859 (1999); M.P. Jacobson, J.P. O'Brien, and R.W. Field, "Anomalous Slow IVR in the $\tilde{X}^1\Sigma_g^+$ State above $10,000\text{ cm}^{-1}$," *J. Chem. Phys.* **109**, 3831 - 3840 (1998); M.P. Jacobson, J.P. O'Brien, R.J. Silbey, and R.W. Field, "Pure Bending Dynamics in the Acetylene $\tilde{X}^1\Sigma_g^+$ State to $15,000\text{ cm}^{-1}$ of Internal Energy," *J. Chem. Phys.* **109**, 121-133 (1998); J.P. O'Brien, M.P. Jacobson, J.J. Sokol, S.L. Coy, and R.W. Field, "Numerical Pattern Recognition in Acetylene Dispersed Fluorescence Spectra," *J. Chem. Phys.* **108**, 7100-7113 (1998); M. Abbouti Temsamani, M. Herman, S.A.B. Solina, J.P. O'Brien, and R.W. Field, "Highly Vibrational Excited $^{12}C_2H_2$ in the $\tilde{X}^1\Sigma_g^+$ State: Complementarity of Absorption and Dispersed Fluorescence Spectra," *J. Chem. Phys.* **105**, 11357-11359 (1997).
- ALT00a. S. Altunata and R. W. Field, "A Statistical Approach to the Study of the Singlet-Triplet Interactions in the 5.5 - 5.7 eV Region of Acetylene," *J. Chem. Phys.* (submitted).
- ALT00b. S. Altunata and R. W. Field, "An Unexpected Application of the Lawrance-Knight Spectral Deconvolution Procedure as a Diagnostic to Retrieve Electronic Coupling Mechanisms Underlying Complex Spectra," in preparation for *J. Chem. Phys.*
- ALT98. S. Altunata, K.L. Cunningham, S. Drucker, C.G. Morgan, and R.W. Field, "Surface Electron Ejection by Laser-Excited Metastable Spectroscopy of C_2H_2 ," Talk TG12, 53rd International Symposium on Molecular Spectroscopy, 1998.
- ALT99. S. Altunata, K.L. Cunningham, S. Drucker, and R.W. Field, "Investigation of the Mechanism of Intersystem Crossing in S_1 Acetylene," Talk TG11, 54th International Symposium on Molecular Spectroscopy.
- ASH87. M.N.R. Ashfold, B. Tutchter, B. Yang, Z.K. Jin, and S.L. Anderson, "Gerade Rydberg States of Acetylene Studied by Multiphoton Ionization and Photoelectron Spectroscopy," *J. Chem. Phys.* **87**, 5105 (1989).
- AVO77. Ph. Avouris, W. M. Gelbart, and M. A. El Sayed, "Nonradiative Electronic Relaxation under Collision-Free Conditions," *Chem. Revs.* **77**, 793-833 (1977).
- BEC97. C. Beck, H. M. Keller, S. Yu Grebenshchikov, and R. Schinke, "Highly Excited Vibrational States of HCP and Their Analysis in Terms of Periodic Orbits: The Genesis of Saddle-Node States and Their Spectroscopic Signature," *J. Chem. Phys.* **107**, 9818-9834 (1997).
- BER76. J. O. Berg, "Extraction of Vibronic Information from Tangled Spectra," *Chem. Phys. Lett.* **41**, 547-551 (1976).
- BIX67. M. Bixon and J. Jortner, "Intramolecular Radiationless Transitions," *J. Chem. Phys.* **48**, 715-726 (1967).
- CHA78. R. Chance, A. Prock, and R. Silbey, "Molecular Fluorescence and Energy Transfer near Interfaces," *Adv. Chem. Phys.* **37**, 1-65 (1978).
- CHE90a. Y.-T. Chen, D. M. Watt, R. W. Field, and K. K. Lehmann, "Observation of Highly Vibrationally Excited $\tilde{X}^1\Sigma^+$ HCP by Stimulated Emission Pumping Spectroscopy," *J. Chem. Phys.* **93**, 2149-2151 (1990).
- COY95. S.L. Coy, D. Chasman, and R.W. Field, "Extracting Dynamical Information from Complex and Congested Spectra: Statistical Pattern Recognition and Parsimonious Trees," pp. 891-942 in *Molecular Dynamics and Spectroscopy by Stimulated Emission Pumping* (H.-L. Dai and R.W. Field, eds.) World Scientific, 1995.

- COY97. S.L. Coy, M.P. Jacobson, and R.W. Field, "Identifying Patterns in Multi-Component Signals by Extended Cross-Correlation," *J. Chem. Phys.* **107**, 8357-8369 (1997).
- CUI96. Q. Cui, K. Morokuma, and J.F. Stanton, "*Ab Initio* MO Studies on the Photodissociation of C_2H_2 from the $S_1^1A_u$ State. Non-Adiabatic Effects and S-T Interaction," *Chem. Phys. Lett.* **263**, 46 (1996).
- CUI98. Q. Cui and K. Morokuma, Personal Communication.
- CUN00a K. L. Cunningham, S. Altunata, M. P. Jacobson, R. Duan, and R. W. Field, "Location and Characterization of the Doorway State in the $3v_3$ Vibrational Level of Acetylene," in preparation for *J. Chem. Phys.*
- CUN00b K. L. Cunningham, M. P. Jacobson, S. Altunata, and R. W. Field, "Extracting Patterns from a SEELEM Spectrum by Extended Auto-Correlation," in preparation for *J. Chem. Phys.*
- DEM75. D. Demoulin, "The shape of Some Excited States of Acetylene", *Chem. Phys.* **11**, 329-341 (1975).
- DOD00 J. A. Dodd, R. B. Lockwood, E. Hwang, S. Miller, and S. J. Lipson, "Formation of OH ($v=0, 1$) by the Reaction of Fast H with O_3 ," *J. Phys. Chem. A* **103**, 7834-7842 (2000).
- DRA94. M. Drabbe, J. Heinze, and W. Leo Meerts, "A Study of the Singlet ~ Triplet Perturbations in the $\tilde{A}-A_u$ State of Acetylene by High Resolution Ultraviolet Spectroscopy", *J. Chem. Phys.* **100**, 165 (1994).
- DRU97. S. Drucker, J.P. O'Brien, P. Patel, and R.W. Field, "The Effects of Triplet Perturbations in Photophysical Processes in the \tilde{A}^1A_u state of Acetylene," *J. Chem. Phys.* **106**, 3423-3426 (1997).
- DUP91. P. Dupré, R. Jost, M. Lombardi, P. G. Green, E. Abramson, and R. W. Field, "Anomalous Behavior of the Anticrossing Density as a Function of Excitation Energy in the C_2H_2 Molecule" *J. Chem. Phys.* **152**, 293-318 (1991).
- DUP93. P. Dupré and P.G. Green, "Characterization of a Large Singlet-Triplet Coupling Matrix Element in the Acetylene Molecule," *Chem. Phys. Lett.* **212**, 555-560 (1993).
- DUP95a. P. Dupré, "Study of Zeeman Anticrossing Spectra of the \tilde{A}^1A_u State of the Acetylene Molecule (C_2H_2) by Fourier Transform: product of $\rho_{vib} \langle V \rangle$ and isomerization barrier," *Chem. Phys.* **196**, 239-266 (1995)
- DUP95b P. Dupré, P.G. Green, and R.W. Field, "Quantum Beat Spectroscopic Studies of Zeeman Anticrossings in the \tilde{A}^1A_u State of the Acetylene (C_2H_2) Molecule," *Chem. Phys.* **196**, 211-238 (1995)
- HAG54. H. Hagstrum, "Theory of Auger Ejection of Electrons from Metals by Ions," *Phys. Rev.* **96**, 336-365 (1954)
- HAG98. F.C. Hagemeister, C.A. Arrington, B.J. Giles, B. Quimpo, L. Zhang, and T.S. Zwier, "Cavity Ringdown Methods for Studying Intramolecular and Intermolecular Dynamics", *J. Phys. Chem.*, **000 000-0000** (1998).
- HCP99. M. Joyeux, D. Sugny, V. Tyng, M.E. Kellman, H. Ishikawa, R.W. Field, C. Beck, and R. Schinke, "Semiclassical Study of the Isomerization States of HCP," *J. Chem. Phys.* **000**, 0000-0000 (2000).; H. Ishikawa, R.W. Field, S.C. Farantos, M. Joyeux, J. Koput, C. Beck, and R. Schinke, "HCP→CPH Isomerization: Caught in the Act!," *Annu. Revs. Phys. Chem.* **50**, 443-484 (1999); H. Ishikawa, C. Nagao, N. Mikami, and Robert W. Field, "Spectroscopic Investigation of the Generation of the 'Isomerization' States: Eigenvector Analysis of the Bend-CP Stretch Polyad," *J. Chem. Phys.* **109**, 492-503 (1998); H. Ishikawa, Chioko Nagao, Naohiko Mikami, and R.W. Field, "Observation of the 'Isomerization States' of HCP by Stimulated Emission Pumping Spectroscopy: Comparison between Theory and Experiment," *J. Chem. Phys.* **106**, 2980-2983 (1997); H. Ishikawa, Y.-T. Chen, Y. Ohshima, B. Rajaram, J. Wang, and R. W. Field, "Stimulated Emission Pumping Spectroscopy of HCP near the Isomerization Barrier: $E_{VIB} \leq 25,315 \text{ cm}^{-1}$," *J. Chem. Phys.* **105**, 7383-7401 (1996).
- HEL87. E. J. Heller, "Quantum Localization and the Rate of Exploration of Phase Space," *Phys. Rev.* **A35**, 1360-1370 (1987).

- HUM97. S.J. Humphrey, C.G. Morgan, A.M. Wodtke, K.L. Cunningham, S. Drucker, and R.W. Field, "Laser Excited Metastable States of Acetylene in the 5.5 - 5.7eV Region," J. Chem. Phys. **107**, 49-53 (1997).
- HUN71 H. E. Hunziker, "A New Technique for Gas-Phase Kinetic Spectroscopy of Molecules in the Triplet State," IBM J. Res. Develop. (1971), 10-26.
- ICL95 A. Kachanov, A. Charvat, and F. Stoeckel, "Intracavity laser spectroscopy with vibronic solid state lasers: I. Spectrotemporal transient behavior of a Ti:sapphire laser," J. Opt. Soc. Amer. B **11**: 2412 - 2421 (1994); A. Kachanov, A. Charvat, and F. Stoeckel, "Intracavity laser spectroscopy with vibronic solid state lasers: II. Influence of the nonlinear mode coupling on the maximum sensitivity of a Ti:sapphire laser," J. Opt. Soc. Amer. B **12**, 970 - 979 (1995); A. Campargue, F. Stoeckel, and M. Chenevier, "High sensitivity intracavity laser absorption spectroscopy: applications to the study of overtone transitions in the visible range," Spectrochimica Acta Rev **13**: 69-88, (1990).
- ING53. C. K. Ingold and G. W. King, "Excited States of Acetylene. Parts I-V," J. Chem. Soc. 2702-2755 (1953).
- INN54. K.K. Innes, "Analysis of the Near Ultraviolet Absorption Spectrum of Acetylene," J. Chem. Phys. **22**, 863 (1954).
- ISH96. H. Ishikawa, Y.-T. Chen, Y. Ohshima, J. Wang, and R. W. Field, "Stimulated Emission Pumping Spectroscopy of HCP near the Isomerization Barrier: $E_{\text{VIB}} \leq 25,315 \text{ cm}^{-1}$," J. Chem. Phys.
- JAC97. M. P. Jacobson, S. L. Coy, and R. W. Field, "Extended Cross-Correlation: A Technique for Spectroscopic Pattern Recognition," J. Chem. Phys. **107**, 8349-8356 (1997).
- JAC98. Matthew P. Jacobson, Stephen L. Coy, Robert W. Field, Steven J. Lipson, Ronald B. Lockwood, David L. Vitito, William A. M. Blumberg and Peter S. Armstrong, "Numerical Pattern Recognition Analysis of CO Atmospheric Simulation Experiments," J. Phys. Chem. **104**, 249-257 (2000).
- JAC99 M. P. Jacobson, "Spectroscopic Patterns Encode Unimolecular Dynamics," Ph. D. Thesis, MIT, 1999.
- KAM70 W. E. Kammer, "Ab Initio SCF and CI Calculations of Linear and Bent Acetylene," Chem. Phys. Lett. **6**, 529-532 (1970)
- KUR98 T. Kurosawa, K. Ohmori, H. Chiba, M. Okunishi, K. Ueda, Y. Sato, A. Z. Devdariani, E. E. Nikitin, "Collision induced absorption in mercury-rare-gas collisions," J. Chem. Phys. **108**, 8101-8109 (1998).
- LAW85. W. D. Lawrance and A. E. W. Knight, "Direct Deconvolution of Extensively Perturbed Spectra: The Singlet-Triplet Molecular Eigenstate Spectrum of Pyrazine," J. Phys. Chem. **89**, 917-925 (1985).
- LAW91. W. D. Lawrance and A. E. W. Knight, "Reply to Kevin Lehmann's Comment," J. Phys. Chem. **95**, 7557-7558 (1991).
- LEF86. H. Lefebvre-Brion and R.W. Field, BOOK: Perturbations in the Spectra of Diatomic Molecules, 465 p., Academic Press, 1986.
- LEH85 K. K. Lehmann, S. C. Ross, and L. L. Lohr, "Experimental and *Ab Initio* Determination of the Bending Potential of HCP," J. Chem. Phys. **82**, 4460-4469 (1985).
- LEH91. K.K. Lehmann, "Comment on 'Direct Deconvolution of Extensively Perturbed Spectra: The Singlet-Triplet Molecular Eigenstate Spectrum of Pyrazine'," J. Phys. Chem. **95**, 7556-7557 (1991).
- LEH96. K. K. Lehmann and D. Romanini, "The Superposition Principle and Cavity Ring-Down Spectroscopy," J. Chem. Phys. **105**, 10263-10277 (1996).
- LIS80. J.M. Lisy and W. Klemperer, "Electric Deflection Studies of Metastable Acetylene," J. Chem. Phys. **72**, 3880-3883 (1980)
- LIS86. H. Lischka, and A. Karpfen, "Ab Initio Calculations on the Excited-States of Pi-Systems. I. Valence Excitations in Acetylene," Chem. Phys. **102**, 77-89 (1986)
- MOR96. C.G. Morgan, M. Drabbels, and A.M. Wodtke, "The Correlated Product State Distribution of Ketene Photodissociation at 308nm," J. Chem. Phys. **104**, 7460-7474 (1996).

- NAK64. T. Nakayama and K. Watanabe, "Absorption and Photoionization Coefficients of Acetylene, Propyne, and 1-Butyne," *J. Chem. Phys.* **40**, 558 (1964).
- NES96. David J. Nesbitt and Robert W. Field, "Vibrational Energy Flow in Highly Excited Molecules: Role of Intramolecular Vibrational Redistribution," *J. Phys. Chem.* **100**, 12,735-12,756 (1996).
- OCH87. N. Ochi and S. Tsuchiya, "Quantum Beat Spectroscopy of Zeeman Splitting and Level Anticrossing of Rotationally Selected Acetylene (\tilde{A}^1A_u 3v₃) under Weak Magnetic Fields", *Chem. Phys. Lett.* **140**, 20 (1987).
- OCH91. N. Ochi, and S. Tsuchiya, "Rovibronic Level Structure of Electronically Excited Acetylene (\tilde{A}^1A_u) in a Supersonic Jet as Studied by Laser-Induced Fluorescence and Zeeman Quantum Beat Spectroscopy," *Chem. Phys.* **152**, 319-336 (1991).
- OHM94. K. Ohmori, T. Kurosawa, H. Chiba, M. Okunishi, and Y. Sato, "Far-Wing Excitation Study on the Fine-Structure Transition of Hg ($^3P_1 - ^3P_0$) in collisions with N₂", *J. Chem. Phys.* **100**, 5381 (1994).
- OHM95. K. Ohmori, T. Kurosawa, H. Chiba, M. Okunishi, K. Ueda, Y. Sato, and E. E. Nikitin, "Far-Wing Excitation Study on the Transit Region of Hg $^3P_1 \rightarrow ^3P_0$ Intra-Multiplet Process in Collisions with N₂ and CO", *J. Chem. Phys.* **102**, 7341 (1995).
- OHM99. K. Ohmori, T. Kurosawa, H. Chiba, M. Okunishi, Y. Sato, A. Z. Devdariani, E. E. Nikitin, "Far-wing excitation study on the transition regions of the metastable mercury atom collisions: Hg(6^3P_2)+N₂ and CO," *Chem. Phys. Lett.* **315**, 411-415 (1999).
- OKA78. Hideo Okabe, *Photochemistry of Small Molecules*, John Wiley and Sons, New York, 1978.
- OKE95. A. O'Keefe and D. A. G. Deacon, "Cavity Ring-Down Optical Spectrometer for Absorption Measurements Using Pulsed Laser Sources," *Rev. Sci. Instrum.* **59**, 2544-2551 (1988); J. J. Scherer, J. B. Paul, A. O'Keefe, and R. J. Saykally, "CRLAS: A New Analytical Technique for Cluster Science," in *Advances in Metal and Semiconductor Clusters*, Vol. 3, pages 149-180 (1995).
- PRA88. J.L. Tomer, K.W. Holtzclaw, and D.W. Pratt, "Phosphorescence Excitation Spectroscopy in Supersonic Jets. The Lowest Triplet-State of Pyrazine," *J. Chem. Phys.* **88**, 1528-1535 (1988); L.H. Spangler and D.W. Pratt, "Laser-Induced Phosphorescence Spectroscopy in Supersonic Jets. The Lowest Triplet-States of Glyoxal, Methylglyoxal, and Biacetyl," *J. Chem. Phys.* **84**, 4789-4796 (1986).
- PRI45. W. C. Price and A. D. Walsh, "The Absorption Spectra of Triple Bond Molecules in the Vacuum Ultra Violet," *Trans. Farad. Soc.* **41**, 381 (1945).
- RUC00. A.F. Ruckstuhl, M. P. Jacobson, R. W. Field, and J. A. Dodd, "Baseline Subtraction Using Robust Local Regression Estimation," *J. Quant. Spectrosc. Radiat. Transf.* **00**, 000-000 (2000).
- SCH86. G.J. Scherer, Y. Chen, R.L. Redington, J.L. Kinsey, and R.W. Field, "An Unsuspected Fermi Perturbation in the Acetylene \tilde{A}^1A_u 3v₃ Level", *J. Chem. Phys.* **85**, 6315-6323 (1986).
- SEK85. K. Seki, "Photochemical Processes of Simple Unsaturated Molecules," Ph.D. Thesis, University of Tokyo, 1985.
- SHE94. C.D. Sherrill and H.F. Schaefer III, "The \tilde{A}^1A " State of Isocyanogen (CNCN)," *J. Chem. Phys.* **100**, 8920 (1994).
- SHE96. C. D. Sherrill, G. Vacek, Y. Yamaguchi, and H. F. Schaefer III, "The \tilde{A}^1A_u State and the T₂ Potential Surface of Acetylene: Implications for Triplet Perturbations in the Fluorescence Spectra of the \tilde{A} -State", *J. Chem. Phys.* **104**, 8507-8515 (1996).
- SNE86a. O. Sne and O. Cheshnovsky, "Surface Ejection of Electrons by Laser-Excited Metastables of Anthracene Derivatives," *Chem. Phys. Lett.* **130**, 53-58 (1986).
- SNE86b. O. Sne and O. Cheshnovsky, "Long Radiative Lifetimes of SO₂ in a Collision-Free Supersonic Molecular Beam," *Chem. Phys. Lett.* **130**, 487-492 (1986).
- SNE88. O. Sne and O. Cheshnovsky, "Triplet Selectivity in Surface Ejection of Electrons by Laser-Excited Metastables of Aniline," *Chem. Phys. Lett.* **146**, 216-220 (1988).

- SNE91. O. Sne and O. Cheshnovsky, "Dynamics of Triplet States in Beam-Isolated Benzaldehyde," *J. Phys. Chem.* **95**, 7154-7164 (1991).
- SOL95. S.A.B. Solina, J. P. O'Brien, R.W. Field, and W.F. Polik, "The Acetylene S_0 Surface: From Dispersed Fluorescence Spectra to Polyads to Dynamics", *Ber. der Bunsengesellschaft f. Phys. Chem.* **99**, 555 (1995).
- SOL96. S.A.B. Solina, J.P. O'Brien, R.W. Field, and W.F. Polik, "Dispersed Fluorescence Spectra of Acetylene Excited via the $\tilde{A}^1A_u \leftarrow \tilde{X}^1\Sigma_g^+$ Origin Band: Recognition of Polyads and Test of Multiresonant Effective Hamiltonian Model for the \tilde{X} -State", *J. Phys. Chem.* **100**, 7797-7809 (1996).
- STA81. P. R. Stannard and W. M. Gelbart, "Intramolecular Vibrational Energy Redistribution," *J. Phys. Chem.* **85**, 3592-3599 (1981).
- STA94. J. F. Stanton, C. Ming-Huang, and P. G. Szalay, "Stationary Points on the S_1 Potential Energy Surface of C_2H_2 ", *J. Chem. Phys.* **101**, 356 (1994).
- SUZ97. T. Suzuki, Y. Shi, and H. Kohguchi, "Detection of Metastable Triplet Acetylene Produced by Intersystem Crossing from the Excited \tilde{A}^1A_u State," *J. Chem. Phys.* **106**, 5292-5295 (1997).
- TAK92. M. Takahashi, M. Fujii, and M. Ito, "Two-Color Double Resonance Spectroscopy via \tilde{A}^1A_u State of Acetylene: 3p Rydberg State and its Renner-Teller Effect," *J. Chem. Phys.* **96**, 6486 (1992).
- UTZ93. A. L. Utz, J. D. Tobiason, E. Carrasquillo M., L. J. Sanders, and F. F. Crim, "The Direct Observation, Assignment, and Partial Deperturbation of the ν_4 and ν_6 Vibrational Fundamentals in \tilde{A}^1A_u Acetylene (C_2H_2)," *J. Chem. Phys.* **98**, 2742-2753 (1993); J. D. Tobiason, A. L. Utz, and F. F. Crim, "The Direct Observation, Assignment, and Partial Deperturbation of ν_5 and $\nu_3+\nu_5$ in \tilde{A}^1A_u Acetylene (C_2H_2)," *J. Chem. Phys.* **99**, 928-936 (1993).
- VAC93. G. Vacek, J. R. Thomas, B. J. DeLeeuw, Y. Yamaguchi, and H. F. Schaefer III, "Isomerization Reactions on the Lowest Potential Energy Hypersurface of Triplet Vinylidene and Triplet Acetylene", *J. Chem. Phys.* **98**, 4766 (1993).
- VAC96. G. Vacek, C.D. Sherrill, Y. Yamaguchi, and H.F. Schaefer, III, "The Anomalous Behavior of the Zeeman Anticrossing Spectra of $\tilde{A}-A_u$ Acetylene: Theoretical Considerations," *J. Chem. Phys.* **104**, 1774-1778 (1996).
- VIT97a. "Infrared Spectral Data Analysis and Remote Sensing Using Pattern Recognition Algorithms" S.J. Lipson, R.B. Lockwood, D.L. Vititoe, C. L. Allred, W.A.M. Blumberg, P.S. Armstrong, M.P. Jacobson, S.L. Coy, and R.W. Field, Presented at the University of Virginia Physics Department Atomic and Molecular Seminar (Charlottesville, VA, April 1997) - Invited
- VIT97b. "Infrared Spectral Data Analysis and Remote Sensing Using Pattern Recognition Algorithms" S.J. Lipson, R.B. Lockwood, D.L. Vititoe, C. L. Allred, W.A.M. Blumberg, P.S. Armstrong, M.P. Jacobson, S.L. Coy, and R.W. Field, Presented at the 1997 Spring Meeting of the American Geophysical Union (Baltimore, MD, May 1997)
- VIT97c. "Disentangling Vibrational Bands in CO Spectra Utilizing Spectroscopic Pattern Recognition Techniques" M.P. Jacobson, S.L. Coy, R.W. Field, S.J. Lipson, R.B. Lockwood, D.L. Vititoe, W.A.M. Blumberg, and P.S. Armstrong, Presented at the 52nd International Symposium on Molecular Spectroscopy (Columbus, OH, June 1997)
- VIT97d. "Infrared Spectral Data Analysis and Remote Sensing Using Pattern Recognition Algorithms" S.J. Lipson, R.B. Lockwood, D.L. Vititoe, C. L. Allred, W.A.M. Blumberg, P.S. Armstrong, M.P. Jacobson, S.L. Coy, and R.W. Field, Presented at the 20th Annual Review Conference of Atmospheric Transmission Models (Hanscom AFB, MA, June 1997)
- VIT97e. "Techniques for Target Detection Using Hyperspectral Data" C. L. Allred, D.L. Vititoe, S.J. Lipson, R.B. Lockwood, L. S. Jeong, W.A.M. Blumberg, P.S. Armstrong, M.P. Jacobson, S.L. Coy, R.W. Field, and W.J. Marinelli, Presented at the Spectral Imaging Technology Workshop (Kirtland AFB, NM, August 1997)

- VIT97f. "Techniques for Target Detection Using Hyperspectral Data" D.L. Vititoe, C. L. Allred, S.J. Lipson, R.B. Lockwood, L. S. Jeong, W.A.M. Blumberg, P.S. Armstrong, M.P. Jacobson, S.L. Coy, and R.W. Field Presented at the Defense Intelligence Agency Advanced Concepts Division 1997 Spectroradiometric Symposium (San Diego, CA, November 1997)
- VIT97g. "Atmospheric Remote Sensing Applications of Spectroscopic Pattern Recognition Algorithms" D.L. Vititoe, S.J. Lipson, C. L. Allred, R.B. Lockwood, W.A.M. Blumberg, P.S. Armstrong, M.P. Jacobson, S.L. Coy, and R.W. Field, Presented at the 1997 Fall Meeting of the American Geophysical Union (San Francisco, CA, December 1997)
- WAT95 J. K. G. Watson, "Franck-Condon Calculations for the $\tilde{A} - \tilde{X}$ Transition of Acetylene," private communication, 1995
- WEN79. H.R. Wendt, H. Hippler, and H.E. Hunziker, "Triplet Acetylene: Near Infrared Electronic Absorption Spectrum of the *cis* Isomer, and Formation from Methylene," J. Chem. Phys. **70**, 4044-4048 (1979)
- WET78. R.W. Wetmore, and H.F. Schaefer, "Triplet Electronic States of Acetylene: *Cis* and *Trans* Structures and Energetics," J. Chem. Phys. **69**, 1648-1654 (1978)
- WOL90. D. Wolff and H. Zacharias, "Lifetimes and Collisional Self-Quenching Rates for $C_2H_2 \tilde{A}^1A_u$ ($v'_3 = 0$ to 4)," Chem. Phys. Lett. **174**, 563 (1990).
- WOO71. R. B. Woodward and R. Hoffman, The Conservation of Orbital Symmetry, Verlag Chemie GmbH/Academic Press, Weinheim, 1971.
- YAM91. K. Yamanouchi, N. Ikeda, S. Tsuchiya, D. M. Jonas, J. K. Lundberg, G. W. Adamson, and R. W. Field, "Vibrationally Highly Excited Acetylene as Studied by Dispersed Fluorescence and Stimulated Emission Spectroscopy: Vibrational Assignment of Feature States", J. Chem. Phys. **95**, 6330 (1991).
- YAM93. Y. Yamaguchi, G. Vacek, and H. F. Schaefer III, "Low-Lying Triplet Electronic States of Acetylene: *cis* 3B_2 and 3A_2 , *trans* 3B_u and 3A_u " Theor. Chim. Acta **86**, 97-113 (1993).
- YAM94. Y. Yamaguchi, G. Vacek, J. R. Thomas, B.J. DeLeeuw, and H. F. Schaefer III, "First and Second Energy Derivative Analyses of the Vinylidene and Acetylene Triplet State Potential Energy Surfaces", J. Chem. Phys. **100**, 4969 (1994).
- ZAL95. P. Zalicki and R. N. Zare, "Cavity Ring-Down Spectroscopy for Quantitative Absorption Measurements," J. Chem. Phys. **102**, 2708-2717 (1995).

III AFOSR Supported Publications (since 1996).

- H. Ishikawa, Y.-T. Chen, Y. Ohshima, B. Rajaram, J. Wang, and R. W. Field, "Stimulated Emission Pumping Spectroscopy of HCP near the Isomerization Barrier: $E_{VIB} \leq 25,315 \text{ cm}^{-1}$ ", J. Chem. Phys. **105**, 7383-7401 (1996).
- R. W. Field, J. P. O'Brien, M. P. Jacobson, S. A. B. Solina, W. F. Polik, and H. Ishikawa, "Intramolecular Dynamics in the Frequency Domain," Adv. in Chem. Phys. **101**, 463-490 (Solvay Conference 1995), 1997.
- S. Drucker, J. P. O'Brien, P. Patel, and R. W. Field, "The Effects of Triplet Perturbers in Photophysical Processes in the A^1A_u state of Acetylene," J. Chem. Phys. **106**, 3423-3426 (1997).
- H. Ishikawa, Chioko Nagao, Naohiko Mikami, and R. W. Field, "Observation of the 'Isomerization States' of HCP by Stimulated Emission Pumping Spectroscopy: Comparison between Theory and Experiment," J. Chem. Phys. **106**, 2980-2983 (1997).
- S. J. Humphrey, C. G. Morgan, A. M. Wodtke, K. L. Cunningham, S. Drucker, and R. Field, "Laser Excited Metastable States of Acetylene in the 5.5-5.7eV Region," J. Chem. Phys. **107**, 49-53(1997).

M. P. Jacobson, S. L. Coy, and R. W. Field, "Extended Cross-Correlation: A Technique for Spectroscopic Pattern Recognition," J. Chem. Phys. 107, 8349-8356 (1997).

S. L. Coy, M. P. Jacobson, and R. W. Field, "Identifying Patterns in Multi-Component Signals by Extended Cross-Correlation," J. Chem. Phys. 107, 8357-8369 (1997).

H. Ishikawa, C. Nagao, N. Mikami, and Robert W. Field, "Spectroscopic Investigation of the Generation of the 'Isomerization' States: Eigenvector Analysis of the Bend-CP Stretch Polyad," J. Chem. Phys. 109, 492-503 (1998).

M. P. Jacobson, S. L. Coy, R. W. Field, S. J. Lipson, R. B. Lockwood, D. L. Vitito, W. A. M. Blumberg, and P. S. Armstrong, "Numerical Pattern Recognition Analysis of CO Atmospheric Simulation Experiments," J. Phys. Chem. 104, 249 - 257 (2000).

H. Ishikawa, R. W. Field, S. C. Farantos, M. Joyeux, J. Koput, C. Beck, and R. Schinke, "HCP→CPH Isomerization: Caught in the Act!," Annu. Revs. Phys. Chem. 50, 443-484 (1999).

M. Joyeux, D. Sugny, V. Tyng, M. E. Kellman, H. Ishikawa, R. W. Field, C. Beck, and R. Schinke, "Semiclassical Study of the Isomerization States of HCP," J. Chem. Phys. 112, 4162-4172 (2000).

A. F. Ruckstuhl, M. P. Jacobson, R. W. Field, and J. A. Dodd, "Baseline Subtraction Using Robust Local Regression Estimation," J. Quant. Spectr. Radiat. Transf. 68, 179 - 193 (2001).

S. Altunata and R. W. Field, "A Statistical Approach to the Study of Singlet-Triplet Interactions in the 5.5 - 5.7 eV Region of Acetylene, J. Chem. Phys. 113, 6640 - 6651 (2000).

IV. Interactions/Transitions: Talks

Talks.

Several talks were given by Robert Field in which the XCC techniques was applied to Dispersed Fluorescence Spectra of C₂H₂. The C₂H₂ DF project is supported by DOE, but the development of XCC was primarily supported by AFOSR .

- | | |
|-------------|--|
| 8/25/96 | American Chemical Society, Invited Talk, "Core-Nonpenetrating Rydberg Series: Spectroscopic Black Holes." |
| 10/24/96 | Optical Society of America, Invited talk, "Now Wrinkles on an Old Plum: Frequency-Modulator Enhanced Magnetic Rotation and Pulsed Frequency Modulation Spectroscopies." |
| 11/6/96 | Discussion Meeting of the Royal Society: Molecular Rydberg Dynamics, Invited talk, "Core-Nonpenetrating Rydberg Series. Spectroscopic Black Holes." |
| 11/18-22/96 | Three talks, Princeton University, Department of Chemistry

"HCP↔HPC Isomerization: Caught in the Act"

"Acetylene Unzipped: Dynamics from Frequency-Domain Dispersed Fluorescence Spectra"

"Spectroscopic Black Holes" |
| 12/3/96 | University of Connecticut, Department of Physics, "Acetylene Unzipped" |

- 1/7/97 Gordon Research Conference, Molecular Energy Transfer, Ventura, CA, "Intramolecular Dynamics in the Frequency Domain"
- 1/16/97 Rowland Institute, "Spectroscopic Black Holes"
- 4/19-5/4/97 Tsinghua University, Beijing, five lectures.
- 6/3/97 Invited Talk, Canadian Society for Chemistry, Windsor, Ontario. "Acetylene Unzipped."
- 6/16-20/97 52nd International Symposium on Molecular Spectroscopy
- RA09 Robert W. Field, "A Dispersed Fluorescence Data Set for the \tilde{X} -State of Acetylene: An Approach Toward Multidimensional Vibrational Spectroscopy."
- RA10 Matthew Jacobson, "Analysis of Acetylene \tilde{X} -State Dynamics Utilizing Spectroscopic Pattern Recognition Techniques."
- RA11 Matthew P. Jacobson, "Disentangling Vibrational Bands in Infrared Spectra of CO by Utilizing Spectroscopic Pattern Recognition Techniques" (with Steven J. Lipson, Ronald B. Lockwood, David L. Vittoe, William A.M. Blumberg, and Peter S. Armstrong)
- RI01 Kevin Cunningham, "Laser Excited Metastable Spectroscopy of C₂H₂"
- RI02 Robert W. Field, "Gateway Mediated Intersystem Crossing in C₂H₂"
- 7/11/97 Laboratoire Photophysique Moléculaire, Orsay, France, "Acetylene Unzipped."
- 10/13/97 CNRS Symposium in honor of Hélène Lefebvre-Brion, Orsay, France. "Perturbations Are Not Accidents," R. W. Field.
- 4/16/98 University of Pennsylvania, "Dispersed Fluorescence Spectra of Acetylene: Order Out of Chaos," R.W. Field.
- 5/1/98 Institute for Theoretical Atomic and Molecular Physics, "Large Amplitude Bending Dynamics of Acetylene," M.P. Jacobson.
- 6/15-19 53rd International Symposium on Molecular Spectroscopy, Ohio State University, Talks MI09, MI10, TB03, TG12, TG13, RF06, RF09, RF10.
- "Mode Beats and the Sensitivity Limit of the Pulsed Cavity Ring-Down Spectroscopy," A.A. Kachanov, S.I. Panov, I. Dubinsky, L.M. Ruslen, M.L. Silva, and R.W. Field.
- "An Optimal Data Acquisition Scheme for a Pulsed Cavity Ring-Down Experiment," A.A. Kachanov, S.I. Panov, I. Dubinsky, L.M. Ruslen, M.L. Silva, and R.W. Field.
- "The Predissociation Mechanism for $2\Sigma^+$ Rydberg States of CaCl," J.O. Clevenger, N.A. Harris, R.W. Field, Jian Li.
- "Surface Electron Ejection by Laser-Excited Metastable Spectroscopy of C₂H₂," S.A. Altunata, K.L. Cunningham, S. Drucker, C.G. Morgan, and R.W. Field.

"Gateway Mediated Intersystem Crossing in C₂H₂," K.L. Cunningham, S. Drucker, S. Altunata, C.G. Morgan, and R.W. Field.

"Spectroscopic Investigation of the Generation of 'Isomerization' States of HCP: Eigenvector Analysis of the Bend-CP Stretch Polyad," H. Ishikawa, C. Nagao, N. Mikami, and R.W. Field.

"Rovibrational Level Structure of Acetylene by Stimulated Emission Pumping: Coriolis Coupling between $\nu_2 + 4\nu_4$ and $7\nu_4$," D.B. Moss, R. Duan, M.P. Jacobson, J.P. O'Brien, and R.W. Field.

"Large Amplitude Bending Dynamics of Acetylene," M.P. Jacobson, J.P. O'Brien, R.J. Silbey and R.W. Field.

8/23-27/98 216th National ACS Meeting, Boston, MA.

Symposium: Special Topics in High-Resolution Spectroscopy: Patterns in Spectra. R.W. Field and K.K. Lehmann co-organized 7 half-day sessions.

Talk #372 "Acetylene Pure Bending Dynamics," M.P. Jacobson, J.P. O'Brien, R.J. Silbey, and R.W. Field.

Talk #376 "Surface Electron Ejection by Laser-Excited Metastable Spectroscopy of Acetylene," K.L. Cunningham, S. Altunata, S. Drucker, C.G. Morgan, and R.W. Field.

Talks by R. Field, K. Cunningham, and S. Altunata at the 52nd (1997), 53rd (1998), 54th (1999), and 55th (2000) "International Symposium on Molecular Spectroscopy (Ohio State University)"

Talk by Selen Altunata at Wesleyan University (2000)

Talks by Robert Field at national ACS meetings (Spring, 2000 and Pacificchem 2000), Edwards Air Force Base (Spring, 2000), International Conference on Photochemistry (San Juan, Puerto Rico, April, 2000), Kobe University (May, 2000), Argonne National Lab (October, 2000), University of California, Riverside (December, 2000).

V. New Discoveries, Inventions, Patents.

None.

VI. Honors

Robert Field: William F. Meggers Prize, Optical Society of America, October 22, 1996
Doctor of Science (honoris causa), Amherst College, June 24, 1997.

Matt Jacobson: Rao Prize, 52nd International Symposium on Molecular Spectroscopy.

Robert Field: Elected to Fellowship in the American Academy of Arts and Sciences, 1998.

Robert Field: Haslam and Dewey Professorship, 1999.

Susan Lang: Dreyfus Undergraduate Research Fellowship, 1997.

Susan Lang: Chemistry Department Prize for Undergraduate Research, 1999.